arXiv:math/9910174v1 [math.AG] 30 Oct 1999

THE HODGE POLYNOMIAL OF $\overline{\mathcal{M}}_{3,1}$

E. GETZLER AND E. LOOIJENGA

Abstract. We give formulas for the Dolbeault numbers of $\overline{\mathcal{M}}_{3,1}$, using the first author's calculations of the weights of the cohomology of $\mathcal{M}_{2,2}$ and $\mathcal{M}_{2,3}$ and the second author's calculation of the weights of the cohomology of $\mathcal{M}_{3,1}$.

Recall that the Serre polynomial of a quasi-projective variety X, or more generally, of a Deligne-Mumford stack whose coarse moduli space is a quasi-projective variety, is the polynomial

$$\mathsf{e}(X) = \sum_{i,j} u^i v^j \sum_{k=0}^{\infty} (-1)^k h^{i,j}(H^k_c(X,\mathbb{C})).$$

If a finite group Γ acts on X, the equivariant Serre polynomial $\mathsf{e}_\Gamma(X)$ is defined in the same way, and takes values in $R(\Gamma)$. For background to all of this, see [1].

If X is smooth and projective, or more generally, if X is a smooth Deligne-Mumford stack whose coarse moduli space is projective, then the Serre polynomial of X equals its Hodge polynomial:

$$\mathsf{e}(X) = \sum_{i,j} (-u)^i (-v)^j h^{i,j}(X).$$

In particular, Poincaré duality implies the functional equation

$$\mathsf{e}(X)(u,v) = q^{\dim(X)}\,\mathsf{e}(X)(u^{-1},v^{-1}),$$

where throughout this paper, q denotes uv. All of this is also true equivariantly.

If $2g-2+n>0$, let $\mathcal{M}_{g,n}$ be the moduli stack of smooth projective curves of genus g together with n distinct marked points, and let $\overline{\mathcal{M}}_{g,n}$ be the moduli stack of stable curves of (arithmetic) genus g with n distinct marked smooth points [4]; these stacks are smooth, of dimension $3g-3+n$, and the coarse moduli space of $\overline{\mathcal{M}}_{g,n}$ is projective. Furthermore, there is a flat map $\overline{\mathcal{M}}_{g,n+1} \to \overline{\mathcal{M}}_{g,n}$ which represents $\overline{\mathcal{M}}_{g,n+1}$ as the universal curve over $\overline{\mathcal{M}}_{g,n}$. In particular, $\overline{\mathcal{M}}_{g,1}$ is the universal stable curve of genus g. In this paper, we calculate $\mathsf{e}(\overline{\mathcal{M}}_{3,1})$. In particular, we show that $h^{i,j}(\overline{\mathcal{M}}_{3,1})=0$ unless $i=j$.

The authors thank the Scuola Normale, Pisa, where this paper was written. The work of the first author is partially funded by the NSF under grant DMS-9704320.

There is a standard identification of $R(\mathbb{S}_n)$ with the space of symmetric functions (in an infinite number of variables $\{x_i \mid i \geq 0\}$) of degree n, which we denote by Λ_n. If X is a quasi-projective variety with action of $\mathbb{S}_n$, let $\mathsf{e}_n(X)$ be its Serre polynomial, thought of as an element of $\Lambda_n[u,v]$.

There is a formula permitting the calculation of the $\mathbb{S}_n$-equivariant Hodge polynomial of $\overline{\mathcal{M}}_{g,n}$ in terms of the $\mathbb{S}_m$-equivariant Serre polynomials of the moduli stacks $\mathcal{M}_{h,m}$, where $h \leq g$ and $2h+m \leq 2g+n$.

Let $\hat{\Lambda}$ be the completion of the ring of symmetric functions:

$$\hat{\Lambda} = \prod_{n=0}^{\infty} \Lambda_n .$$

We may identify $\hat{\Lambda}$ with the algebra of power series in the variables

$$h_n = \sum_{i_1 \leq \dots \leq i_n} x_{i_1} \dots x_{i_n} .$$

Introduce an auxilliary variable λ, and let $\hat{\Lambda}[\![\lambda]\!]$ be the ring of power series in λ with coefficients in the ring of symmetric functions. We may identify $\hat{\Lambda} \otimes \mathbb{Q}$ with the algebra of power series in the variables

$$p_n = \sum_{i=0}^{\infty} x_i^n .$$

The ring $\hat{\Lambda}[\![\lambda]\!]$ is a (special) lambda-ring: the Adams operations on $\hat{\Lambda}[\![\lambda]\!]$ are determined by the formulas $\psi_k(p_n) = p_{kn}$ and $\psi_k(\lambda) = \lambda^k$. Let Exp be the operation

$$\mathsf{Exp}(f) = \sum_{k=0}^{\infty} \sigma_k(f) = \exp\left(\sum_{k=0}^{\infty} \frac{\psi_k(f)}{k} \right) : \lambda\hat{\Lambda}[\![\lambda]\!] \to 1 + \lambda\hat{\Lambda}[\![\lambda]\!],$$

and let Log be its inverse, given by the formula

$$\mathsf{Log}(f) = \sum_{k=0}^{\infty} \frac{\mu(k)}{k} \psi_k(f) : 1 + \lambda\hat{\Lambda}[\![\lambda]\!] \to \lambda\hat{\Lambda}[\![\lambda]\!].$$

Let Δ be the differential operator on $\hat{\Lambda}$ (and by extension of coefficients, on $\hat{\Lambda}[\![\lambda]\!]$) give by the formula

$$\Delta = \sum_{k=1}^{\infty} \lambda^{2k} \left(\frac{k}{2} \frac{\partial^2}{\partial p_k^2} + \frac{\partial}{\partial p_{2k}} \right) .$$

We may define two elements of $\hat{\Lambda}[\![\lambda]\!]$, by the formulas

$$\begin{aligned} \Psi &= \sum_{2g-2+n>0, n>0} \lambda^{2g-2+n} \, \mathsf{e}_{\mathbb{S}_n}(\overline{\mathcal{M}}_{g,n}) \quad \text{and} \\ \Phi &= \sum_{2g-2+n>0, n>0} \lambda^{2g-2+n} \, \mathsf{e}_{\mathbb{S}_n}(\overline{\mathcal{M}}_{g,n}). \end{aligned}$$

The following is Theorem (8.13) of [3].

Theorem 1. $\Psi = \mathsf{Log}\big(\exp(\Delta)\,\mathsf{Exp}(\Phi)\big)$

We see that to use this formula to calculate $\mathsf{e}(\overline{\mathcal{M}}_{3,1})$, we need the Serre polynomial of $\mathcal{M}_{3,1}$, together with the following equivariant Serre polynomials:

$\mathcal{M}_{g,n}$	$\mathsf{e}_{\mathbb{S}_n}(\mathcal{M}_{g,n})$
$\mathcal{M}_{0,3}$	s_3
$\mathcal{M}_{0,4}$	$qs_4 - s_{2^2}$
$\mathcal{M}_{0,5}$	$q^2 s_5 - qs_{32} + s_{31^2}$
$\mathcal{M}_{0,6}$	$q^3 s_6 - q^2 s_{42} + q(s_{41^2} + s_{321}) - (s_{41^2} + s_{3^2} + s_{2^2 1^2})$
$\mathcal{M}_{0,7}$	$q^4 s_7 - q^3 s_{52} + q^2(s_{3^2 1} + s_{51^2} + s_{421})$
	$- q(s_{51^2} + s_{43} + s_{421} + s_{2^3 1} + s_{41^3} + s_{3^2 1} + s_{321^2})$
	$+ s_{321^2} + s_{52} + s_{31^4} + s_{32^2} + s_{421}$
$\mathcal{M}_{1,1}$	qs_1
$\mathcal{M}_{1,2}$	$q^2 s_2$
$\mathcal{M}_{1,3}$	$q^3 s_3 - s_{1^3}$
$\mathcal{M}_{1,4}$	$q^4 s_4 - q^2 s_4 - qs_{21^2} + s_{31}$
$\mathcal{M}_{1,5}$	$q^5 s_5 - q^3(s_5 + s_{41}) + q^2(s_{32} - s_{31^2})$
	$+ q(s_{41} + s_{32} + s_{31^2}) - (s_5 + s_{32} + s_{2^2 1} + s_{1^5})$
$\mathcal{M}_{2,1}$	$q^4 s_1 + q^3 s_1$
$\mathcal{M}_{2,2}$	$q^5 s_2 + q^4(s_2 + s_{1^2}) - s_2$
$\mathcal{M}_{2,3}$	$q^6 s_3 + q^5(s_3 + s_{21}) - q^4 s_3 - q(s_3 + s_{21})$

Here, if $\mu = (\mu_1 \geq \cdots \geq \mu_\ell)$ is a partition of n, s_μ denotes the associated Schur polynomial, given by the Jacobi-Trudi formula $s_\mu = \det(h_{\mu_i+j-i})$. The above Serre polynomials are calculated in [1] (for genus 0 and 1) and [2] (for genus 2).

Applying Theorem 1 (and taking advantage of J. Stembridge's symmetric function package `SF` [6] for `maple`), we see that

$$\mathsf{e}(\overline{\mathcal{M}}_{3,1}) = \mathsf{e}(\mathcal{M}_{3,1}) + 3q^6 + 15q^5 + 29q^4 + 29q^3 + 16q^2 + 4q.$$

Theorem 2. *The Hodge polynomial* $\mathsf{e}(\overline{\mathcal{M}}_{3,1})$ *of* $\overline{\mathcal{M}}_{3,1}$ *equals*

$$q^7 + 5q^6 + 16q^5 + 29q^4 + 29q^3 + 16q^2 + 5q + 1.$$

Proof. We must show that $\mathsf{e}(\mathcal{M}_{3,1}) = q^7 + 2q^6 + q^5 + q + 1$. This almost, but not quite, follows from [5]. There, it was shown that the polynomial

$$\sum_{i,j,k} u^i v^j t^k h^{i,j}(H^k_c(\mathcal{M}_{3,1},\mathbb{C})) = q^7 t^{14} + q^6 2t^{12} + q^5 t^{10} + q(t^8 + t^7) + 2t^6,$$

which would imply that $\mathsf{e}(\mathcal{M}_{3,1}) = q^7 + 2q^6 + q^5 + 2$.

Unfortunately, there was a small error in that calculation; namely, two terms were omitted from the calculation (4.8) of the Poincaré polynomial of

$W\backslash T'$ in case R is of type E_7, associated to the Dynkin subdiagrams $D_6 \times A_1$ and $D_4 \times A_1^2$ of E_7. Once they are taken into account (the first contributes t^7, the second qt^8), we see that

$$\begin{aligned}\sum_{i,j,k} u^i v^j t^k h^{i,j}(H^k(\mathcal{M}_{3,1},\mathbb{C})) = q^7t^{14} + q^6 2t^{12} + q^5t^{10} + qt^8 + t^7 \\ + (aqt+b)(t^7+t^6),\end{aligned}$$

where a and b equal 0 or 1. This proves the theorem. ☐

References

[1] E. Getzler, *Resolving mixed Hodge structures of configuration spaces.* Duke Math. J. **96** (1999), 175–203.

[2] E. Getzler, *Topological recursion relations in genus* 2, in "Integrable systems and algebraic geometry (Kobe/Kyoto, 1997)," World Sci. Publishing, River Edge, NJ, 1998, pp. 73–106.

[3] E. Getzler and M. Kapranov, *Modular operads,* Compositio Math. **110** (1998), 65–126.

[4] F.F. Knudsen, *The projectivity of the moduli space of stable curves II. The stacks* $\overline{\mathcal{M}}_{g,n}$*,* Math. Scand. **52** (1983), 161–189.

[5] E. Looijenga, *Cohomology of* $\mathcal{M}_3$ *and* $\mathcal{M}_3^1$*,* in "Mapping class groups and moduli spaces of Riemann surfaces (Göttingen, 1991/Seattle, WA, 1991)," Contemp. Math. **150**, Amer. Math. Soc., Providence, RI, 1993, pp. 205–228.

[6] J. Stembridge, *A Maple package for symmetric functions.* `http://www.math.lsa.umich.edu/~jrs/maple.html`.

Northwestern University, 2033 Sheridan Rd., Evanston, IL 60208-2730, USA
E-mail address: `getzler@math.nwu.edu`

Utrecht University, P.O. Box 80010, 3508 TA Utrecht, The Netherlands
E-mail address: `looijeng@math.uu.nl`